

UNCLASSIFIED

AD 299 487

*Reproduced
by the*

ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

63-3-1

299 487

MEMORANDUM
RM-3482-PR
MARCH 1963

CATALOGED BY ASTIA
AC AD 100

299 487

A NOTE ON DIFFERENTIAL APPROXIMATION AND ORTHOGONAL POLYNOMIALS

Richard Bellman

ASTIA
APR 2 1963
TEA A

PREPARED FOR:
UNITED STATES AIR FORCE PROJECT RAND

The **RAND** Corporation
SANTA MONICA • CALIFORNIA

MEMORANDUM

RM-3482-PR

MARCH 1963

**A NOTE ON
DIFFERENTIAL APPROXIMATION
AND ORTHOGONAL POLYNOMIALS**

Richard Bellman

This research is sponsored by the United States Air Force under Project RAND—contract No. AF 49(638)-700 monitored by the Directorate of Development Planning, Deputy Chief of Staff, Research and Development, Hq USAF. Views or conclusions contained in this Memorandum should not be interpreted as representing the official opinion or policy of the United States Air Force.

The **RAND** *Corporation*

1700 MAIN ST • SANTA MONICA • CALIFORNIA

PREFACE

Part of the Project RAND research program consists of basic supporting studies. The mathematical research presented here is concerned with an analytic technique for approximating a given function by a sum of exponentials. The results that are obtained have applications in control theory, circuit synthesis, and numerical analysis.

SUMMARY

We consider a simple analytic technique for finding linear differential equations which are approximately satisfied by a given function, and thus obtain approximations to a given function by a sum of exponentials.

CONTENTS

PREFACE.	11
SUMMARY.111
Section	
1. INTRODUCTION	1
2. FOURIER TRANSFORMS	2
3. ORTHOGONAL POLYNOMIALS AND MINIMIZATION.	3
4. STABILITY.	3
REFERENCES	5

A NOTE ON DIFFERENTIAL APPROXIMATION AND ORTHOGONAL POLYNOMIALS

1. INTRODUCTION

The problem of obtaining an exponential polynomial of the form $\sum_{i=1}^N b_i e^{\lambda_i t}$ which closely approximates a given function $f(t)$ in an interval $a \leq t \leq b$ is a problem of some difficulty if we allow both the coefficients and the exponents to be unknowns. Either of the criteria of fit,

$$(1.1) \quad \max_{a \leq t \leq b} \left| f(t) - \sum_{i=1}^N b_i e^{\lambda_i t} \right|,$$

or

$$(1.2) \quad \int_a^b \left| f(t) - \sum_{i=1}^N b_i e^{\lambda_i t} \right|^2 dt$$

lead to difficulties; see Lanczos [1].

In this note, we wish to consider a different way of measuring the closeness of $f(t)$ to a sum of exponentials. If $f(t)$ were a function of the form $\sum_{i=1}^N b_i e^{\lambda_i t}$, it would satisfy a linear differential equation of the form

$$(1.3) \quad f^{(N)} + c_1 f^{(N-1)} + \dots + c_N f = 0.$$

Hence, let us attempt to determine real coefficients c_1 which minimize the integral

$$(1.4) \quad \int_{-\infty}^{\infty} (f^{(N)} + c_1 f^{(N-1)} + \dots + c_N f)^2 dt.$$

We call this differential approximation.

Results of the type we obtain have applications in the fields of control theory, circuit synthesis, and numerical analysis.

2. FOURIER TRANSFORMS

It is clear that we must impose some conditions on the function $f(t)$ in order to pose the problem. Assume then that $f^{(k)} \in L^2(-\infty, \infty)$ for $k = 0, 1, 2, \dots, N$. Then if

$$(2.1) \quad \begin{aligned} g(s) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(t) e^{ist} dt, \\ f(t) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(s) e^{-ist} ds, \end{aligned}$$

we have

$$(2.2) \quad f^{(k)}(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (-is)^k g(s) e^{-ist} ds.$$

Using the Plancherel-Parseval formula, we have

$$(2.3) \quad \begin{aligned} &\int_{-\infty}^{\infty} (f^{(N)} + c_1 f^{(N-1)} + \dots + c_N f)^2 dt \\ &= \int_{-\infty}^{\infty} |g(s)|^2 |c_N - is c_{N-1} + (-is)^2 c_{N-2} + \dots|^2 ds \\ &= \int_{-\infty}^{\infty} |g(s)|^2 [(c_N - s^2 c_{N-2} + \dots)^2 + s^2 (c_{N-1} - s^2 c_{N-3} + \dots)^2] ds. \end{aligned}$$

3. ORTHOGONAL POLYNOMIALS AND MINIMIZATION

It is clear that a change of variable reduces the problem to that of finding the orthogonal polynomials associated respectively with the weight functions $|g(s)|^2$ and $s^2|g(s)|^2$. Since these orthogonal polynomials can be constructed in a systematic fashion, we have a simple way of obtaining the coefficients c_1 , and further measures of the asymptotic behavior as $N \rightarrow \infty$; see Szego [2].

A direct treatment of the question of minimizing the quadratic form of (1.4) would not be simple computationally for large N (although a Schmidt orthogonalization could be used), and would not readily furnish asymptotic behavior. The technique presented above is most useful in connection with the treatment of various classical functions where $g(s)$ has a simple analytic form.

Similar results can be obtained for the case where the interval is finite or semi-infinite, but not of the same simplicity.

4. STABILITY

The study of the precise connection between the solution of the linear differential equation

$$(4.1) \quad u^{(N)} + c_1 u^{(N-1)} + \dots + c_N u = 0,$$

~~4~~

and the function $f(t)$ leads to a stability question
which we shall study elsewhere.

REFERENCES

1. Lanczos, C., Applied Analysis, Prentice Hall, Englewood Cliffs, New Jersey, 1956.
2. Szego, G., Orthogonal Polynomials, American Mathematical Society Colloquium Publications, Vol. 23, New York, 1939.

See also

Laning, J. H., and R. H. Battin, Random Processes in Automatic Control, McGraw-Hill Book Co., Inc., New York, 1956.

Jaroch, O., "Approximation by Exponential Functions," Aplikace Mat., Vol. 7, 1962, pp. 249-264.